



AMERICAN SOCIETY OF
PLASTIC SURGEONS®

Informed Consent

Augmentation Mammoplasty with Silicone Gel-Filled Implants and Mastopexy

Informed Consent – Augmentation Mammoplasty with Silicone Gel-Filled Implants/Mastopexy

INSTRUCTIONS

This is an informed-consent document that has been prepared to help inform you about augmentation mammoplasty and mastopexy surgery with silicone gel-filled implants, its risks, as well as alternative treatment(s).

It is important that you read this information carefully and completely. Please initial each page, indicating that you have read the page and sign the consent for surgery as proposed by your plastic surgeon and agreed upon by you.

GENERAL INFORMATION

In November, 2006, silicone gel-filled breast implant devices were approved by the United States Food and Drug Administration (FDA) for use in breast augmentation and reconstruction.

Augmentation mammoplasty is a surgical operation performed to enlarge the female breasts for a number of reasons:

- To enhance the body contour of a woman, who for personal reasons feels that her breast size is too small.
- To correct a loss in breast volume after pregnancy.
- To balance breast size, when there exists a significant difference between the size of the breasts.
- To restore breast shape after partial or total loss of the breasts for various conditions.
- To correct a failure for breast development due to a severe breast abnormality.
- To correct or improve results of existing breast implants for cosmetic or reconstructive reasons.

Breast implant surgery is contraindicated in women with untreated breast cancer or pre-malignant breast disorders, active infection anywhere in the body, or individuals who are currently pregnant or nursing.

Individuals with a weakened immune system (currently receiving chemotherapy or drugs to suppress the immune system), conditions that interfere with blood clotting or wound healing, or have reduced blood supply to the breast tissue from prior surgery or radiation therapy treatments may be at greater risk for complications and poor surgical outcome.

Silicone breast implants are approved by the FDA for use in women that are at least 22 years of age. Women that meet this age criteria may utilize the silicone implants for cosmetic breast augmentation or for revision surgery to correct or improve results of earlier cosmetic breast augmentation. There is no age restriction on breast reconstruction procedures to restore breast shape after cancer, trauma, or severe breast abnormalities.

Breast enlargement is accomplished by inserting a breast implant either behind the breast tissue, or partially or completely under the chest muscles. Incisions are made to keep scars as inconspicuous as possible, usually under the breast, around a portion of areola, or in the armpit. According to the FDA it is not recommended to use the peri-umbilical approach to insert gel-filled implants. Breast implants may be manufactured in a variety of shapes, sizes, and with either smooth or textured surfaces. The method of implant selection and size, along with surgical approach for inserting and positioning breast implants will depend on your preferences, your anatomy and your surgeon's recommendation. The shape and size of the breasts prior to surgery will influence both the recommended treatment and the final results. If the breasts are not the same size or shape before surgery, it is unlikely that they will be completely symmetrical afterward.

Conditions which involve sagging of the breast or diminished skin tone (stretch marks) may require additional surgical procedures (breast lift) to reposition the nipple and areola upward and to remove loose skin.

Patients undergoing augmentation mammoplasty surgery must consider the following:

- Breast augmentation or reconstruction with silicone gel-filled implants may not be a onetime surgery.
- Breast implants of any type are not considered lifetime devices. They cannot be expected to last forever. You will likely require future surgery for implant replacement or removal.
- Changes that occur to the breasts following augmentation or reconstruction with implants are not reversible. There may be an unacceptable appearance to the breast if you later choose to have breast implants removed.
- Large volume primary augmentation or revision with larger sized implants in excess of dimensional planning for your chest and breast size may increase the risk of complications such as implant extrusion, hematoma, infection, palpable implant folds, and visible skin wrinkling requiring surgical intervention to correct these complications.

ALTERNATIVE TREATMENTS

Augmentation mammoplasty with silicone gel-filled implants is an elective surgical operation. Alternative treatment would consist of not undergoing the surgical procedure or use of external breast prostheses or padding, saline-filled implants, or the transfer of other body tissues to enlarge/rebuild breast size. Risks and potential complications are associated with alternative surgical forms of treatment.

INHERENT RISKS OF AUGMENTATION MAMMAPLASTY SURGERY

Every surgical procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications or adverse events associated with them. In addition, every procedure has limitations in terms of the outcome that patients will achieve afterwards. Additional information concerning breast implants may be obtained from the FDA, package-insert sheets supplied by the implant manufacturer, or other information pamphlets required by individual state laws.

An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. While all patients do not experience these complications or adverse events, you should discuss each of them with your plastic surgeon to make sure you understand all possible consequences of breast augmentation. Adverse events associated with breast implants can be inherent to this type of implanted medical device or relate to complications of a surgical procedure. Additional advisory information regarding this subject should be reviewed by patients considering surgery that involves breast implants.

While every patient experiences her own individual risks and benefits following breast implant surgery, clinical data suggests that most women will be satisfied with the outcome of breast implant surgery despite the occurrence of problems inherent with the surgery.

SPECIFIC RISKS OF SILICONE GEL-FILLED BREAST IMPLANTS

Implants:

Breast implants, similar to other medical devices, can fail. When a silicone gel-filled implant ruptures, the gel material is usually contained within the scar tissue surrounding the implant (intracapsular rupture). In some cases, the gel may escape beyond the capsule layer and go into the breast tissue itself (extracapsular rupture and gel migration) or to more distant locations. Migrated silicone gel may be difficult or impossible to remove. Rupture of a breast implant may or may not produce local firmness in the breast. Patients are advised to refer to individual manufacturer's informational materials regarding the incidence of device rupture reported during pre-market studies.

It is impossible to predict the biologic response that a patient's tissues will exhibit to the placement of breast implants or how you will heal following surgery.

Rupture can occur as a result of an injury, from no apparent cause or during mammography. Rupture of a silicone breast implant is most often undetected (silent rupture). It is possible to damage an implant at the time of surgery. Damaged or broken implants cannot be repaired. According to the FDA, ruptured or damaged implants require replacement or removal. Breast implants can wear out, they are not guaranteed to last a lifetime and future surgery may be required to replace one or both implants.

A MRI (magnetic resonance imaging) study is advised to evaluate the possibility of implant rupture, yet it may not be 100% accurate in diagnosing implant integrity. It should be noted that the FDA recommends regular screening MRI examinations. Specifically, patients are advised to follow recommendations for serial MRI examinations, starting at 3 years after surgery and then every 2 years thereafter.

Capsular Contracture:

Scar tissue, which forms internally around the breast implant, can tighten and make the breast round, firm, and possibly painful. Excessive firmness of the breasts can occur soon after surgery or years later. The occurrence of symptomatic capsular contracture is not predictable. The incidence of symptomatic capsular contracture can be expected to increase over time. Capsular contracture may occur on one side, both sides or not at all. It is more common with implant placement in front of the chest muscle layer. Treatment for capsular contracture may require surgery, implant replacement, or implant removal. Capsular contracture may reoccur after surgical procedures to treat this condition and it occurs more often in revision augmentation than primary augmentation. Some surgeons believe that preventative antibiotics during dental work and treatment for sinus infections and urinary tract infections may decrease this incidence. Discuss this with your surgeon. There may be an Off-Label FDA use for a drug called Singulair, which may have a softening effect on the capsule.

Implant Extrusion / Tissue Necrosis:

Lack of adequate tissue coverage or infection may result in exposure and extrusion of the implant through the skin. Tissue breakdown (necrosis) has been reported with the use of steroid drugs, after chemotherapy/radiation to breast tissue, due to smoking, microwave diathermy, and excessive heat or cold therapy. In some cases, incision sites fail to heal normally. Atrophy of breast tissue may occur. An implant may become visible at the surface of the breast as a result of the device pushing through layers of skin. If tissue break down occurs and the implant becomes exposed, implant removal may be necessary. Permanent scar deformity may occur.

Skin Wrinkling and Rippling:

Visible and palpable wrinkling of implants and breast skin can occur. Some wrinkling is normal and expected with silicone gel-filled breast implants. This may be more pronounced in patients who have silicone gel-filled implants with textured surfaces or thin breast tissue. Palpable wrinkling and/or folds may be confused with palpable tumors and questionable cases must be investigated.

Calcification:

Calcium deposits can form in the scar tissue surrounding the implant and may cause pain, firmness, and be visible on mammography. These deposits must be identified as different from calcium deposits that are a sign of breast cancer. Should this occur, additional surgery may be necessary to remove and examine calcifications.

Chest Wall Irregularities:

Chest wall irregularities have been reported secondary to the use of tissue expanders and breast implants. Residual skin irregularities at the ends of the incisions or “dog ears” are always a possibility when there is excessive redundant skin. This may improve with time, or it can be surgically corrected.

Implant Displacement and Tissue Stretching:

Displacement, rotation, or migration of a breast implant may occur from its initial placement and can be accompanied by discomfort and/or distortion in breast shape (visible rippling of the skin). Unusual techniques of implant placement may increase the risk of displacement or migration. Additional surgery may be necessary to attempt to correct this problem. It may not be possible to resolve this problem once it has occurred.

Surface Contamination of Implants:

Skin oil, lint from surgical drapes, or talc may become deposited on the surface of the implant at the time of insertion. The consequences of this are unknown.

Unusual Activities and Occupations:

Activities and occupations which have the potential for trauma to the breast could potentially break or damage breast implants, or cause bleeding/seroma.

Silicone Gel Bleed:

The evidence is mixed regarding whether there are any clinical consequences associated with silicone gel bleed. Over time, extremely small amounts of silicone gel material and platinum can pass through the shell layer of the implant and coat the outside of the implant. Studies indicate that a small amounts of platinum in its most biologically compatible (zero oxidation) state are contained within silicone gel. Microgram amounts of platinum in this state have been found to diffuse outside of breast implants. This may contribute to capsular contracture and lymph node swelling. The overall body of available evidence supports that the extremely low levels of gel bleed is of no clinical consequence.

Change in Nipple and Skin Sensation:

You may experience a diminished (or loss of) sensitivity of the nipples and the skin of your breast. After 6 months, 85 percent of patients have normal sensation. Partial or permanent loss of nipple and skin sensation may occur occasionally. Changes in sensation may affect sexual response or the ability to breast feed a baby.

Anaplastic Large Cell Lymphoma (ALCL):

Women with saline and silicone gel breast implants may have a very small and possibly increased risk of developing anaplastic large cell lymphoma (ALCL) in the scar capsule adjacent to the implant. This is a very rare disease and is currently being investigated as to its relationship to breast implants, and whether this is even a cancer or a Lymphoproliferative Disorder. ALCL is an extremely rare cancer of the immune system which can occur anywhere in the body. The National Cancer Institute estimated 1 in 500,000 women per year in the U.S. are diagnosed with ALCL. ALCL in the breast is even rarer with approximately 1 in 32,000 women in U.S diagnosed per year. The relationship between breast implants and ALCL is unclear and is currently under investigation. In most cases, women observed changes in the look or feel of the area surrounding the implant after their initial surgical sites were fully healed.

Patients with breast implants should be followed by a surgeon over time and seek professional care for implant-related symptoms such as pain, lumps, swelling, or asymmetry. Patients should monitor their breast implants with routine breast self exams and follow standard medical recommendations for imaging (e.g. Mammography, Ultrasound, MRI). Abnormal screening results or implant-related symptoms may result in additional costs and expenses for tests and/or procedures to properly diagnose and treat your condition. Tests and procedures could include but may not be limited to: obtaining breast fluid or tissue for pathology and laboratory evaluation and surgery to remove the scar capsule around the breast implant, implant removal or implant replacement.

Breast Disease:

Current medical information does not demonstrate an increased risk of breast cancer in women who have breast implant surgery for either cosmetic or reconstructive purposes. Individuals with a personal history or family history of breast cancer may be at a higher risk of developing breast cancer than a woman with no family history of this disease. It is recommended that all women perform periodic self-examination of their breasts, have mammography according to American Cancer Society guidelines, and seek professional care should a breast lump be detected. In the event that suspicious tissue is identified prior to or during breast surgery, additional tests and therapy with corresponding expenses may be warranted.

SPECIFIC RISKS OF BREAST LIFT (MASTOPEXY) SURGERY

GENERAL INFORMATION

Breast lift or mastopexy is a surgical procedure to raise and reshape sagging breasts. Factors such as pregnancy, nursing, weight change, aging and gravity produce changes in the appearance of a woman's breasts. As the skin loses its elasticity, the breasts often lose their shape and begin to sag. Breast lift or mastopexy is a surgery performed by plastic surgeons to raise and reshape sagging breasts. This operation can also reduce the size of the areola, the darker skin around the nipple. If your breasts are small or have lost volume after pregnancy, breast implants inserted in conjunction with mastopexy can increase both firmness and size. The best candidates for mastopexy are healthy, emotionally stable women who have realistic expectations about what this type of surgery can accomplish. Breasts of any size can be lifted, but the results may not last as long in women with heavy, large breasts. Mastopexy does leave permanent, noticeable scars on the breasts, and the breasts will be smaller than your current size. There are a variety of different surgical techniques used for the reshaping and lifting of the female breast.

ALTERNATIVE TREATMENTS

Mastopexy is an elective surgical operation. Alternative treatment would consist of not undergoing the surgical procedure or wearing supportive undergarments to lift sagging breasts. If breasts are large and sagging, a reduction mammoplasty may be considered. Risks and potential complications are also associated with alternative surgical forms of treatment.

INHERENT RISKS OF BREAST LIFT (MASTOPEXY) SURGERY

Every surgical procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications associated with them. In addition, every procedure has limitations. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with your plastic surgeon to make sure you understand all possible consequences of breast lift (mastopexy) surgery.

Change in Nipple and Skin Sensation:

You may experience a diminished (or loss of) sensitivity of the nipples and the skin of your breast. Partial or permanent loss of nipple and skin sensation can occur after a mastopexy in one or both nipples. Changes in sensation may affect sexual response or the ability to breast feed a baby.

Breast Augmentation and Simultaneous Mastopexy:

Patients who choose to undergo breast implant augmentation and elect to have it at the same time as a breast lift (mastopexy) may be at increased risk of necrosis of skin, nipples, and breast tissue due to decreased blood supply to the tissues.

Mastopexy Performed at the Time of Breast Implant Removal Surgery:

Patients who choose to undergo simultaneous removal of breast implants and capsules and elect to have it at the same time as a breast lift (mastopexy) may be at increased risk of necrosis of skin, nipples, and breast tissue due to decreased blood supply to the tissues from earlier surgery. Risks associated with the removal of breast implants are covered in a separate informed-consent document.

Use of Acellular Dermal Matrix:

In order to perform the breast lift surgery, your plastic surgeon may choose to use biological materials. Most commonly, these materials are derived from human cadaver skin or pig skin. These materials are generally processed and do not carry any viable cells. You should ask your surgeon about these materials. They provide help to support the breast tissues and become populated with your cells, becoming similar to your own tissue. These acellular products may produce fluid and require drains for a prolonged period of time.

Skin Contour Irregularities:

Contour and shape irregularities may occur after mastopexy. Visible and palpable wrinkling may occur. One breast may be smaller than the other. Nipple position and shape will not be identical one side to the next. Residual skin irregularities at the ends of the incisions or “dog ears” are always a possibility when there is excessive redundant skin. This may improve with time, or it can be surgically corrected.

Delayed Healing:

Wound disruption or delayed wound healing is possible. Some areas of the breast skin or nipple region may not heal normally and may take a long time to heal. Areas of skin or nipple tissue may die. This may require frequent dressing changes or further surgery to remove the non-healed tissue. Individuals who have decreased blood supply to breast tissue from past surgery or radiation therapy may be at increased risk for delayed wound healing and poor surgical outcome. Smokers have a greater risk of skin loss and wound healing complications.

Breast Disease:

Breast disease and breast cancer can occur independently of breast lift surgery. Individuals with a personal history or family history of breast cancer may be at a higher risk of developing breast cancer than a woman with no family history of this disease. It is recommended that all women perform periodic self-examination of their breasts, have mammography according to American Cancer Society guidelines, and seek professional care should a breast lump be detected. In the event that suspicious tissue is identified prior to or during breast surgery, additional tests and therapy with corresponding expenses may be warranted.

GENERAL RISKS OF SURGERY

Healing Issues:

Certain medical conditions, dietary supplements and medications may delay and interfere with healing. Patients with massive weight loss may have a healing delay that could result in the incisions coming apart, infection, and tissue changes resulting in the need for additional medical care, surgery, and prolonged hospitalizations. Patients with diabetes or those taking medications such as steroids on an extended basis may have prolonged healing issues. Smoking will cause a delay in the healing process, often resulting in the need for additional surgery. There are general risks associated with healing such as swelling, bleeding, possibility of additional surgery, prolonged recovery, color changes, shape changes, infection, not meeting patient goals and expectations, and added expense to the patient. There may also be a longer recovery due to the length of surgery and anesthesia.

Bleeding:

It is possible, though unusual, to experience a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require emergency treatment to drain accumulated blood or you may require a blood transfusion, though such occurrences are rare. Increased activity too soon after surgery can lead to increased chance of bleeding and additional surgery. It is important to follow postoperative instructions and limit exercise and strenuous activity for the instructed time. Do not take any aspirin or anti-inflammatory medications for at least ten days before or after surgery, as this may increase the risk of bleeding. Non-prescription “herbs” and dietary supplements can increase the risk of surgical bleeding. Hematoma can occur at any time, usually in the first three weeks following injury to the operative area. If blood transfusions are necessary to treat blood loss, there is the risk of blood-related infections such as hepatitis and HIV (AIDS). Heparin medications that are used to prevent blood clots in veins can produce bleeding and decreased blood platelets.

In breast implant surgery, hematoma may contribute to capsular contracture, infection or other problems.

Infection:

Infection is unusual after surgery. Should an infection occur, additional treatment including antibiotics, hospitalization, or additional surgery may be necessary. It is important to tell your surgeon of any other infections, such as ingrown toenail, insect bite, or urinary tract infection. Remote infections, infection in other part of the body, may lead to an infection in the operated area.

Infection in Breast Implant Patients:

Subacute or chronic infections may be difficult to diagnose. Should an infection occur, treatment including antibiotics, possible removal of the implant, or additional surgery may be necessary. Infections with the presence of a breast implant are harder to treat than infections in normal body tissues. If an infection does not respond to antibiotics, the breast implant may have to be removed. After the infection is treated, a new breast implant can usually be reinserted. It is rare that an infection would occur around an implant from a bacterial infection elsewhere in the body, however, prophylactic antibiotics may be considered for subsequent dental or other surgical procedures. In extremely rare instances, life-threatening infections, including toxic shock syndrome have been noted after breast implant surgery. Individuals with an active infection in their body should not undergo surgery, including breast augmentation. Although infection is unusual after this type of surgery, it may appear in the immediate post-operative period or at any time following the insertion of a breast implant. It is important to tell your surgeon of any other infections, such as ingrown toenail, insect bite, or urinary tract infection. Remote infections, infection in other part of the body, may lead to an infection in the operated area.

Scarring:

All surgery leaves scars, some more visible than others. Although good wound healing after a surgical procedure is expected, abnormal scars may occur within the skin and deeper tissues. Scars may be unattractive and of different color than the surrounding skin tone. Scar appearance may also vary within the same scar. Scars may be asymmetrical (appear different on the right and left side of the body). There is the possibility of visible marks in the skin from sutures. In some cases scars may require surgical revision or treatment.

Firmness:

Excessive firmness can occur after surgery due to internal scarring. The occurrence of this is not predictable. Additional treatment including surgery may be necessary.

Change in Skin Sensation:

It is common to experience diminished (or loss of) skin sensation in areas that have had surgery. Diminished (or complete loss of) skin sensation may not totally resolve.

Skin Contour Irregularities:

Contour and shape irregularities may occur. Visible and palpable wrinkling of skin may occur. Residual skin irregularities at the ends of the incisions or “dog ears” are always a possibility when there is excessive redundant skin. This may improve with time, or it can be surgically corrected.

Skin Discoloration / Swelling:

Some bruising and swelling will normally occur. The skin in or near the surgical site can appear either lighter or darker than surrounding skin. Although uncommon, swelling and skin discoloration may persist for long periods of time and, in rare situations, may be permanent.

Informed Consent – Augmentation Mammoplasty with Silicone Gel-Filled Implants/Mastopexy

Skin Sensitivity:

Itching, tenderness, or exaggerated responses to hot or cold temperatures may occur after surgery. Usually this resolves during healing, but in rare situations it may be chronic.

Major Wound Separation:

Wounds may separate after surgery. Should this occur, additional treatment including surgery may be necessary.

Sutures:

Most surgical techniques use deep sutures. You may notice these sutures after your surgery. Sutures may spontaneously poke through the skin, become visible or produce irritation that requires suture removal.

Delayed Healing:

Wound disruption or delayed wound healing is possible. Some areas of the skin may not heal normally and may take a long time to heal. Areas of skin may die. This may require frequent dressing changes or further surgery to remove the non-healed tissue. Individuals who have decreased blood supply to tissue from past surgery or radiation therapy may be at increased risk for wound healing and poor surgical outcome. Smokers have a greater risk of skin loss and wound healing complications.

Damage to Deeper Structures:

There is the potential for injury to deeper structures including nerves, blood vessels, muscles, and lungs (pneumothorax) during any surgical procedure. The potential for this to occur varies according to the type of procedure being performed. Injury to deeper structures may be temporary or permanent.

Fat Necrosis:

Fatty tissue found deep in the skin might die. This may produce areas of firmness within the skin. Additional surgery to remove areas of fat necrosis may be necessary. There is the possibility of contour irregularities in the skin that may result from fat necrosis.

Seroma:

Infrequently, fluid may accumulate between the skin and the underlying tissues following surgery, trauma or vigorous exercise. Should this problem occur, it may require additional procedures for drainage of fluid. Excess fluid accumulation around an implant secondary to too much activity too early may increase capsular contracture occurrence.

Surgical Anesthesia:

Both local and general anesthesia involves risk. There is the possibility of complications, injury, and even death from all forms of surgical anesthesia or sedation.

Shock:

In rare circumstances, your surgical procedure can cause severe trauma, particularly when multiple or extensive procedures are performed. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. If surgical shock occurs, hospitalization and additional treatment would be necessary.

Pain:

You will experience pain after your surgery. Pain of varying intensity and duration may occur and persist after surgery. Chronic pain may occur very infrequently from nerves becoming trapped in scar tissue or due to tissue stretching.

Cardiac and Pulmonary Complications:

Pulmonary complications may occur secondary to blood clots (pulmonary emboli), fat deposits (fat emboli) or partial collapse of the lungs after general anesthesia. Pulmonary emboli can be life-threatening or fatal in some circumstances. Inactivity and other conditions may increase the incidence of blood clots traveling to the lungs causing a major blood clot that may result in death. It is important to discuss with your physician any past history of swelling in your legs or blood clots that may contribute to this condition. Cardiac complications are a risk with any surgery and anesthesia, even in patients without symptoms. If you experience shortness of breath, chest pains, or unusual heart beats, seek medical attention immediately. Should any of these complications occur, you may require hospitalization and additional treatment.

Informed Consent – Augmentation Mammoplasty with Silicone Gel-Filled Implants/Mastopexy

Venous Thrombosis and Sequelae:

Thrombosed veins, which resemble cords, occasionally develop in the area of the breast or around IV sites, and usually resolve without medical or surgical treatment. It is important to discuss with your surgeon any birth control pills you are taking. Certain high estrogen pills may increase your risk of thrombosed veins.

Allergic Reactions:

In rare cases, local allergies to tape, suture material and glues, blood products, topical preparations or injected agents have been reported. Serious systemic reactions including shock (anaphylaxis) may occur in response to drugs used during surgery and prescription medicines. Allergic reactions may require additional treatment.

Drug Reactions:

Unexpected drug allergies, lack of proper response to medication, or illness caused by the prescribed drug are possibilities. It is important for you to inform your physician of any problems you have had with any medication or allergies to medication, prescribed or over the counter, as well as medications you now regularly take.

Asymmetry:

Symmetrical body appearance may not result after surgery. Factors such as skin tone, fatty deposits, skeletal prominence, and muscle tone may contribute to normal asymmetry in body features. Most patients have differences between the right and left side of their bodies before any surgery is performed. Additional surgery may be necessary to attempt to diminish asymmetry.

Surgical Wetting Solutions:

There is the possibility that large volumes of fluid containing dilute local anesthetic drugs and epinephrine that is injected into fatty deposits during surgery may contribute to fluid overload or systemic reaction to these medications. Additional treatment including hospitalization may be necessary.

Persistent Swelling (Lymphedema):

Persistent swelling can occur following surgery.

Unsatisfactory Result:

Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. The body is not asymmetric and almost everyone has some degree of unevenness which may not be recognized in advance. One side of the face may be slightly larger, one side of the face droopier. The breast and trunk area exhibits the same possibilities. Many of such issues cannot be fully corrected with surgery. The more realistic your expectations as to results, the better your results will be in your eye. Some patients never achieve their desired goals or results, at no fault of the surgeon or surgery. You may be disappointed with the results of surgery. Asymmetry, unanticipated shape and size, loss of function, wound disruption, poor healing, and loss of sensation may occur after surgery. Size may be incorrect. Unsatisfactory surgical scar location or appearance may occur. It may be necessary to perform additional surgery to improve your results.

ADDITIONAL ADVISORIES

Smoking, Second-Hand Smoke Exposure, Nicotine Products (Patch, Gum, Nasal Spray):

Patients who are currently smoking or use tobacco or nicotine products (patch, gum, or nasal spray) are at a greater risk for significant surgical complications of skin dying and delayed healing and additional scarring. Individuals exposed to second-hand smoke are also at potential risk for similar complications attributable to nicotine exposure. Additionally, smoking may have a significant negative effect on anesthesia and recovery from anesthesia, with coughing and possibly increased bleeding. Individuals who are not exposed to tobacco smoke or nicotine-containing products have a significantly lower risk of this type of complication. Please indicate your current status regarding these items below:

___ I am a non-smoker and do not use nicotine products. I understand the potential risk of second-hand smoke exposure causing surgical complications.

___ I am a smoker or use tobacco / nicotine products. I understand the risk of surgical complications due to smoking or use of nicotine products.

___ I have smoked and stopped approximately _____ ago. I understand I may still have the effects and therefore risks from smoking in my system, if not enough time has lapsed.

Informed Consent – Augmentation Mammoplasty with Silicone Gel-Filled Implants/Mastopexy

It is important to refrain from smoking at least 6 weeks before surgery and until your physician states it is safe to return, if desired. I acknowledge that I will inform my physician if I continue to smoke within this time frame, and understand that for my safety, the surgery, if possible, may be delayed.

Smoking may have such a negative effect on your surgery that a urine test just before surgery may be done which will prove the presence of Nicotine. If positive, your surgery may be cancelled and your surgery, scheduling fee, and other prepaid amounts may be forfeited. Honestly disclose smoking to your surgeon.

Sleep Apnea / CPAP:

Individuals who have breathing disorders such as “Obstructive Sleep Apnea” and who may rely upon CPAP devices (constant positive airway pressure) or utilize nighttime oxygen are advised that they are at a substantive risk for respiratory arrest and death when they take narcotic pain medications following surgery. This is an important consideration when evaluating the safety of surgical procedures in terms of very serious complications, including death, that relate to pre-existing medical conditions. Surgery may be considered only with monitoring afterwards in a hospital setting in order to reduce risk of potential respiratory complications and to safely manage pain following surgery.

Please consider the following symptoms of sleep apnea:

- I am frequently tired upon waking and throughout the day
- I have trouble staying asleep at night
- I have been told that I snore or stop breathing during sleep
- I wake up throughout the night or constantly turn from side to side
- I have been told that my legs or arms jerk while I'm sleeping
- I make abrupt snorting noises during sleep
- I feel tired or fall asleep during the day

It is important for you to inform and discuss any of the above symptoms that you have experienced with your surgeon.

Medications and Herbal Dietary Supplements:

There are potential adverse reactions that occur as the result of taking over-the-counter, herbal, and/or prescription medications. Aspirin and medications that contain aspirin interfere with bleeding. These include non-steroidal anti-inflammatories such as Motrin, Advil, and Aleve. It is very important not to stop drugs that interfere with platelets, such as Plavix, which is used after a stent. It is important if you have had a stent and are taking Plavix that you inform the plastic surgeon. Stopping Plavix may result in a heart attack, stroke and even death. Be sure to check with your physician about any drug interactions that may exist with medications which you are already taking. If you have an adverse reaction, stop the drugs immediately and call your plastic surgeon for further instructions. If the reaction is severe, go immediately to the nearest emergency room. When taking the prescribed pain medications after surgery, realize that they can affect your thought process and coordination. Do not drive, do not operate complex equipment, do not make any important decisions and do not drink any alcohol while taking these medications. Be sure to take your prescribed medication only as directed.

Sun Exposure – Direct or Tanning Salon:

The effects of the sun are damaging to the skin. Exposing the treated areas to sun may result in increased scarring, color changes, and poor healing. Patients who tan, either outdoors or in a salon, should inform their surgeon and either delay treatment, or avoid tanning until the surgeon says it is safe to resume. The damaging effect of sun exposure occurs even with the use sun block or clothing coverage.

Travel Plans:

Any surgery holds the risk of complications that may delay healing and your return to normal life. Please let the surgeon know of any travel plans, important commitments already scheduled or planned, or time demands that are important to you, so that appropriate timing of surgery can occur. There are no guarantees that you will be able to resume all activities in the desired time frame.

Long-Term Results:

Subsequent alterations in the appearance of your body may occur as the result of aging, sun exposure, weight loss, weight gain, pregnancy, menopause or other circumstances not related to your surgery.

Interference with Sentinel Lymph Node Mapping Procedures:

Breast surgery procedures that involve cutting through breast tissue, similar to a breast biopsy, can potentially interfere with diagnostic procedures to determine lymph node drainage of breast tissue to stage breast cancer.

Body-Piercing Procedures:

Individuals who currently wear body-piercing jewelry in the surgical region are advised that an infection could develop from this activity. Piercing may be done 6 months post-surgery, but is **NOT** recommended.

Future Pregnancy and Breast Feeding:

This surgery is not known to interfere with pregnancy. If you are planning a pregnancy, your breast skin may stretch and offset the results of surgery. You may have more difficulty breast feeding after this operation.

Female Patient Information:

It is important to inform your plastic surgeon if you use birth control pills, estrogen replacement, or if you suspect you may be pregnant. Many medications including antibiotics may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy. Please consult with your OBGYN to develop a plan to discontinue your hormone therapy for up to 2 weeks prior to surgery.

Intimate Relations After Surgery:

Surgery involves coagulating of blood vessels and increased activity of any kind may open these vessels leading to a bleed, or hematoma. Activity that increases your pulse or heart rate may cause additional bruising, swelling, and the need for return to surgery to control bleeding. It is wise to refrain from intimate physical activities until your physician states it is safe.

Mental Health Disorders and Elective Surgery:

It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvement rather than perfection. Complications or less than satisfactory results are sometimes unavoidable, may require additional surgery and often are stressful. Please openly discuss with your surgeon, prior to surgery, any history that you may have of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately predicted.

DVT/PE Risks and Advisory:

There is a risk of blood clots, Deep Vein Thrombosis (DVT) and Pulmonary Embolus (PE) with every surgical procedure. It varies with the risk factors below. The higher the risk factors, the greater the risk and the more involved you must be in both understanding these risks and, when permitted by your physician, walking and moving your legs. There may also be leg stockings, squeezing active leg devices, and possibly medicines to help lower your risk. There are many conditions that may increase or affect risks of clotting. Inform your doctor about any past or present history of any of the following:

- _____ Past History of Blood Clots
- _____ Family History of Blood Clots
- _____ Birth Control Pills
- _____ Swollen Legs
- _____ History of Cancer
- _____ Large Dose Vitamins
- _____ Varicose Veins
- _____ Past Illnesses of the Heart, Liver, Lung, or Gastrointestinal Tract.
- _____ I understand the risks relating to DVT/PE and how important it is to comply with therapy as discussed with my surgeon. The methods of preventative therapy include:
 - _____ Early ambulation when allowed
 - _____ Compression devices (SCD/ICD)
 - _____ ASA protocol when allowed (Aspirin)
 - _____ Heparin protocol when allowed
 - _____ Enoxaparin protocol when allowed

The risks of DVT/PE may be almost as great as the prophylactic therapy when involving Aspirin, Heparin, and Exoxaparin. Be aware that if your surgery is elective, those patients with very high risks should consider not proceeding with such elective surgery.

ADDITIONAL SURGERY NECESSARY (Re-Operations)

There are many variable conditions that may influence the long-term result of surgery. It is unknown how your tissue may respond or how wound healing will occur after surgery. Secondary surgery may be necessary to perform additional tightening or repositioning of body structures. Should complications occur, additional surgery or other treatments may be necessary. Even though risks and complications occur infrequently, the risks cited are particularly associated with this surgery. Other complications and risks can occur but are even more uncommon. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. In some situations, it may not be possible to achieve optimal results with a single surgical procedure. You and your surgeon will discuss the options available should additional surgery be advised. There may be additional costs and expenses for such additional procedures, including surgical fees, facility and anesthesia fees, pathology and lab testing.

PATIENT COMPLIANCE

Follow all physician instructions carefully; this is essential for the success of your outcome. It is important that the surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the time of healing. Personal and vocational activity needs to be restricted. Protective dressings and drains should not be removed unless instructed by your plastic surgeon. Successful post-operative function depends on both surgery and subsequent care. Physical activity that increases your pulse or heart rate may cause bruising, swelling, fluid accumulation and the need for return to surgery. It is wise to refrain from intimate physical activities after surgery until your physician states it is safe. It is important that you participate in follow-up care, return for aftercare, and promote your recovery after surgery.

REVISION POLICY

Surgical revision surgery is a common part of elective surgery. Your procedure will not stop you from aging, sagging, scarring, or experiencing ongoing skin changes that are more genetically controlled. If revision surgery is either desired or advisable within one year after the initial surgery, there may be a physician's fee. Additionally, there may be fees associated with the hospital, facility, anesthesia, pathology, lab, and any supplies such as implants, etc. Revision policy and courtesy discounts only apply to patients who comply with post-op orders and visits.

REGULATORY MATTERS

According to FDA regulations, you must comply with the submission of personal information to a device registry before surgery and afterwards if breast implants are used as a component of your breast surgery.

HEALTH INSURANCE

Most health insurance companies exclude coverage for cosmetic surgical operations or any resulting complications. Please carefully review your health insurance subscriber-information pamphlet. Most insurance plans exclude coverage for secondary or revisionary surgery due to complications of cosmetic surgery. It is unethical and fraudulent to bill insurance for cosmetic procedures. We cannot participate in such activities.

FINANCIAL RESPONSIBILITIES

The cost of surgery involves several charges for the services provided. The total includes fees charged by your surgeon, the cost of surgical supplies, anesthesia, laboratory tests, and possible outpatient hospital charges, depending on where the surgery is performed. Depending on whether the cost of surgery is covered by an insurance plan, you will be responsible for necessary co-payments, deductibles, and charges not covered. The fees charged for this procedure do not include any potential future costs for additional procedures that you elect to have or require in order to revise, optimize, or complete your outcome. Additional costs may occur should complications develop from the surgery. Secondary surgery or hospital day-surgery charges involved with revision surgery will also be your responsibility. In signing the consent for this surgery/procedure, you acknowledge that you have been informed about its risk and consequences and accept responsibility for the clinical decisions that were made along with the financial costs of all future treatments.

___ I understand that with cosmetic surgery, I am responsible for the surgical fees quoted to me, as well as additional fees for anesthesia, facility (OR), and possibly laboratory, X-ray, and pathology fees.
Surgicenters, Outpatient Centers and Hospitals often have rules that certain tissue /implants removed during surgery must be sent for evaluation that may result in additional fees. Please check with your surgeon for approximate additional costs you will be responsible for.

___ I understand that there will be a non-refundable fee for booking and scheduling your surgery of \$ _____, which is a part of the overall surgical fee.

Informed Consent – Augmentation Mammoplasty with Silicone Gel-Filled Implants/Mastopexy

DISCLAIMER

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s), including no surgery. The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed-consent documents should not be considered all-inclusive in defining other methods of care and risks encountered. Your plastic surgeon may provide you with additional or different information which is based on all the facts in your particular case and the current state of medical knowledge.

Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.

CONSENT FOR SURGERY / PROCEDURE or TREATMENT

1. I hereby authorize Dr. Poulter and such assistants as may be selected to perform the following procedure or treatment: Augmentation Mammoplasty with Silicone Gel-Filled Implants and mastopexy

I have received the following information sheet: augmentation Mammoplasty with Silicone Gel-Filled Implants and mastopexy

2. I recognize that during the course of the operation and medical treatment or anesthesia, unforeseen conditions may necessitate different procedures than those above. I therefore authorize the above physician and assistants or designees to perform such other procedures that are in the exercise of his or her professional judgment necessary and desirable. The authority granted under this paragraph shall include all conditions that require treatment and are not known to my physician at the time the procedure is begun.
3. I consent to the administration of such anesthetics considered necessary or advisable. I understand that all forms of anesthesia involve risk and the possibility of complications, injury, and sometimes death.
4. I understand what my surgeon can and cannot do, and understand there are no warranties or guarantees, implied or specific about my outcome. I have had the opportunity to explain my goals and understand which desired outcomes are realistic and which are not. All of my questions have been answered, and I understand the inherent (specific) risks to the procedures I seek, as well as those additional risks and complications, benefits, and alternatives. Understanding all of this, I elect to proceed.
5. I consent to be photographed or televised before, during, and after the operation(s) or procedure(s) to be performed, including appropriate portions of my body, for medical, scientific or educational purposes, provided my identity is not revealed by the pictures.
6. For purposes of advancing medical education, I consent to the admittance of observers to the operating room.
7. I consent to the disposal of any tissue, medical devices or body parts that may be removed.
8. I consent to the utilization of blood products should they be deemed necessary by my surgeon and/or his/her appointees, and I am aware that there are potential significant risks to my health with their utilization.
9. I authorize the release of my Social Security number to appropriate agencies for legal reporting and medical-device registration. Social security number: _____ - _____ - _____
10. I understand that the surgeons' fees are separate from the anesthesia and hospital charges, and the fees are agreeable to me. If a secondary procedure is necessary, further expenditure will be required.
11. I realize that not having the operation is an option.
12. IT HAS BEEN EXPLAINED TO ME IN A WAY THAT I UNDERSTAND:
- a. THE ABOVE TREATMENT OR PROCEDURE TO BE UNDERTAKEN
 - b. THERE MAY BE ALTERNATIVE PROCEDURES OR METHODS OF TREATMENT
 - c. THERE ARE RISKS TO THE PROCEDURE OR TREATMENT PROPOSED

I CONSENT TO THE TREATMENT OR PROCEDURE AND THE ABOVE LISTED ITEMS (1-12).
I AM SATISFIED WITH THE EXPLANATION.

Patient or Person Authorized to Sign for Patient

Date _____ Witness _____